

Lucas Taylor

Publications in the area of Research with the L3 Experiment at CERN

- [1] P. Achard et al., Study of resonance formation in the mass region 1400-MeV to 1500-MeV through the reaction $\gamma\gamma \rightarrow K_0(S) K^{\pm} \pi^{\pm}$, *JHEP* **03**, 018 (2007).
- [2] P. Achard et al., Study of inclusive strange-baryon production and search for pentaquarks in two-photon collisions at LEP, *Eur. Phys. J.* **C49**, 395–410 (2007).
- [3] P. Achard et al., A search for flaring very-high-energy cosmic gamma-ray sources with the L3+C muon spectrometer, *Astropart. Phys.* **25**, 298–310 (2006).
- [4] P. Achard et al., Analysis of the $\pi^+ \pi^- \pi^+ \pi^-$ and $\pi^+ \pi^0 \pi^- \pi^0$ final states in quasi-real two-photon collisions at LEP, *Phys. Lett.* **B638**, 128–139 (2006).
- [5] P. Achard et al., Measurement of hadron and lepton-pair production in $e^+ e^-$ collisions at $s^{*(1/2)} = 192\text{-GeV} - 208\text{-GeV}$ at LEP, *Eur. Phys. J.* **C47**, 1–19 (2006).
- [6] S. Schael et al., Search for neutral MSSM Higgs bosons at LEP, *Eur. Phys. J.* **C47**, 547–587 (2006).
- [7] P. Achard et al., Measurement of the mass and the width of the W boson at LEP, *Eur. Phys. J.* **C45**, 569–587 (2006).
- [8] P. Achard et al., Measurement of the running of the electromagnetic coupling at large momentum-transfer at LEP, *Phys. Lett.* **B623**, 26–36 (2005).
- [9] P. Achard et al., Compton scattering of quasi-real virtual photons at LEP, *Phys. Lett.* **B616**, 145–158 (2005).
- [10] P. Achard et al., Neutral-current four-fermion production in $e^+ e^-$ interactions at LEP, *Phys. Lett.* **B616**, 159–173 (2005).
- [11] P. Achard et al., Measurement of the shadowing of high-energy cosmic rays by the moon: A search for TeV-energy antiprotons, *Astropart. Phys.* **23**, 411–434 (2005).

- [12] P. Achard et al., Measurement of exclusive $\rho^+ \rho^-$ production in mid- virtuality two-photon interactions and study of the $\gamma \gamma^* \rightarrow \rho \rho$ process at LEP, Phys. Lett. **B615**, 19–30 (2005).
- [13] P. Achard et al., Z-boson production with two unobserved, back-to-back, hard photons at LEP, Phys. Lett. **B613**, 118–127 (2005).
- [14] P. Achard et al., Measurement of the cross section for open-beauty production in photon photon collisions at LEP, Phys. Lett. **B619**, 71–81 (2005).
- [15] P. Achard et al., Measurement of the photon structure function $F_2(\gamma)$ with the L3 detector at LEP, Phys. Lett. **B622**, 249–264 (2005).
- [16] P. Achard et al., Search for an invisibly-decaying Higgs boson at LEP, Phys. Lett. **B609**, 35–48 (2005).
- [17] P. Achard et al., Inclusive jet production in two-photon collisions at LEP, Phys. Lett. **B602**, 157–166 (2004).
- [18] P. Achard et al., Measurement of exclusive $\rho^0 \rho^0$ production in mid- virtuality two-photon interactions at LEP, Phys. Lett. **B604**, 48–60 (2004).
- [19] P. Achard et al., Measurement of the atmospheric muon spectrum from 20-GeV to 3000-GeV, Phys. Lett. **B598**, 15–32 (2004).
- [20] P. Achard et al., Studies of hadronic event structure in $e^+ e^-$ annihilation from 30-GeV to 209-GeV with the L3 detector, Phys. Rept. **399**, 71–174 (2004).
- [21] P. Achard et al., Measurement of the cross section of W-boson pair production at LEP, Phys. Lett. **B600**, 22–40 (2004).
- [22] P. Achard et al., Study of the $e^+ e^- \rightarrow Z \gamma$ process at LEP and limits on triple neutral-gauge-boson couplings, Phys. Lett. **B597**, 119–130 (2004).
- [23] P. Achard et al., Search for branons at LEP, Phys. Lett. **B597**, 145–154 (2004).
- [24] P. Achard et al., Search for anomalous couplings in the Higgs sector at LEP, Phys. Lett. **B589**, 89–102 (2004).
- [25] P. Achard et al., Measurement of exclusive $\rho^+ \rho^-$ production in high- Q^2 two-photon collisions at LEP, Phys. Lett. **B597**, 26–38 (2004).
- [26] P. Achard et al., Measurement of triple gauge boson couplings of the W boson at LEP, Phys. Lett. **B586**, 151–166 (2004).
- [27] P. Achard et al., Muon pair and tau pair production in two-photon collisions at LEP, Phys. Lett. **B585**, 53–62 (2004).
- [28] P. Achard et al., Inclusive Λ production in two-photon collisions at LEP, Phys. Lett. **B586**, 140–150 (2004).

- [29] P. Achard et al., Measurement of the Z-boson mass using $e^+ e^-$ to Z gamma events at centre-of-mass energies above the Z pole, Phys. Lett. **B585**, 42–52 (2004).
- [30] P. Achard et al., Study of spin and decay-plane correlations of W bosons in the $e^+ e^-$ to $W^+ W^-$ process at LEP, Eur. Phys. J. **C40**, 333–341 (2005).
- [31] P. Achard et al., Single- and multi-photon events with missing energy in $e^+ e^-$ collisions at LEP, Phys. Lett. **B587**, 16–32 (2004).
- [32] P. Achard et al., Flavour independent search for neutral Higgs bosons at LEP, Phys. Lett. **B583**, 14–27 (2004).
- [33] P. Achard et al., Search for doubly-charged Higgs bosons at LEP, Phys. Lett. **B576**, 18–28 (2003).
- [34] P. Achard et al., Search for charged Higgs bosons at LEP, Phys. Lett. **B575**, 208–220 (2003).
- [35] P. Achard et al., Search for a Higgs boson decaying to weak boson pairs at LEP, Phys. Lett. **B534**, 28–38 (2002).
- [36] P. Achard et al., Search for scalar leptons and scalar quarks at LEP, Phys. Lett. **B580**, 37–49 (2004).
- [37] P. Achard et al., Search for colour singlet and colour reconnection effects in hadronic Z decays at LEP, Phys. Lett. **B581**, 19–30 (2004).
- [38] P. Achard et al., Z boson pair-production at LEP, Phys. Lett. **B572**, 133–144 (2003).
- [39] P. Achard et al., Measurement of exclusive $\rho^0 \rho^0$ production in two-photon collisions at high Q^2 at LEP, Phys. Lett. **B568**, 11–22 (2003).
- [40] P. Achard et al., Measurement of branching fractions of tau hadronic decays, CERN-EP-2003-019.
- [41] P. Achard et al., Search for colour reconnection effects in $e^+ e^-$ to $W^+ W^-$ to hadrons through particle-flow studies at LEP, Phys. Lett. **B561**, 202–212 (2003).
- [42] R. Barate et al., Search for the standard model Higgs boson at LEP, Phys. Lett. **B565**, 61–75 (2003).
- [43] P. Achard et al., Search for excited leptons at LEP, Phys. Lett. **B568**, 23–34 (2003).
- [44] P. Achard et al., Proton antiproton pair production in two-photon collisions at LEP, Phys. Lett. **B571**, 11–20 (2003).
- [45] P. Achard et al., Measurement of W polarisation at LEP, Phys. Lett. **B557**, 147–156 (2003).
- [46] P. Achard et al., Study of the $e^+ e^-$ to Z $e^+ e^-$ process at LEP, Phys. Lett. **B561**, 73–81 (2003).
- [47] P. Achard et al., Inclusive charged hadron production in two-photon collisions at LEP, Phys. Lett. **B554**, 105–114 (2003).
- [48] P. Achard et al., Search for neutral Higgs bosons of the minimal supersymmetric standard model in $e^+ e^-$ interactions at \sqrt{s} up to 209-GeV, Phys. Lett. **B545**, 30–42 (2002).

- [49] P. Achard et al., Production of single W bosons at LEP and measurement of W W gamma gauge coupling parameters, Phys. Lett. **B547**, 151–163 (2002).
- [50] P. Achard et al., Search for single top production at LEP, Phys. Lett. **B549**, 290–300 (2002).
- [51] P. Achard et al., Measurement of Bose-Einstein correlations in $e^+ e^- \rightarrow W^+ W^-$ events at LEP, Phys. Lett. **B547**, 139–150 (2002).
- [52] P. Achard et al., The $e^+ e^- \rightarrow Z \gamma \gamma \rightarrow q \bar{q} \gamma \gamma$ reaction at LEP and constraints on anomalous quartic gauge boson couplings, Phys. Lett. **B540**, 43–51 (2002).
- [53] P. Achard et al., Measurement of genuine three-particle Bose-Einstein correlations in hadronic Z decay, Phys. Lett. **B540**, 185–198 (2002).
- [54] P. Achard et al., Search for a Higgs boson decaying into two photons at LEP, Phys. Lett. **B534**, 28–38 (2002).
- [55] P. Achard et al., Lambda and Sigma0 pair production in two photon collisions at LEP, Phys. Lett. **B536**, 24–33 (2002).
- [56] P. Achard et al., Inclusive D^{*+} production in two-photon collisions at LEP, Phys. Lett. **B535**, 59–69 (2002).
- [57] P. Achard et al., Determination of α_S from Hadronic Event Shapes in e^+e^- Annihilation at $192 < \sqrt{s} < 208$ GeV, Phys. Lett. **B536**, 217–228 (2002).
- [58] P. Achard et al., Study of multiphoton final states and tests of QED in $e^+ e^-$ collisions at \sqrt{s} up to 209-GeV, Phys. Lett. **B531**, 28–38 (2002).
- [59] P. Achard et al., Study of the $W^+ W^- \gamma$ process and limits on anomalous quartic gauge boson couplings at LEP, Phys. Lett. **B527**, 29–38 (2002).
- [60] P. Achard et al., Search for R-parity violating decays of supersymmetric particles in $e^+ e^-$ collisions at LEP, Phys. Lett. **B524**, 65–80 (2002).
- [61] P. Achard et al., Measurement of the charged-particle multiplicity and inclusive momentum distributions in Z decays at LEP, (2001).
- [62] P. Achard et al., $f_1(1285)$ formation in two photon collisions at LEP, Phys. Lett. **B526**, 269–277 (2002).
- [63] P. Achard et al., Double-tag events in two-photon collisions at LEP, Phys. Lett. **B531**, 39–51 (2002).
- [64] P. Achard et al., Inclusive π^0 and $K_0(S)$ production in two-photon collisions at LEP, Phys. Lett. **B524**, 44–54 (2002).
- [65] P. Achard et al., Bose-Einstein correlations of neutral and charged pions in hadronic Z decays, Phys. Lett. **B524**, 55–64 (2002).
- [66] P. Achard et al., Search for heavy isosinglet neutrino in $e^+ e^-$ annihilation at LEP, Phys. Lett. **B517**, 67–74 (2001).

- [67] P. Achard et al., Standard model Higgs boson with the L3 experiment at LEP, Phys. Lett. **B517**, 319–331 (2001).
- [68] P. Achard et al., Measurement of the topological branching fractions of the tau lepton at LEP, Phys. Lett. **B519**, 189–198 (2001).
- [69] P. Achard et al., Search for heavy neutral and charged leptons in $e^+ e^-$ annihilation at LEP, Phys. Lett. **B517**, 75–85 (2001).
- [70] M. Acciarri et al., Measurement of the tau branching fractions into leptons, Phys. Lett. **B507**, 47–60 (2001).
- [71] M. Acciarri et al., Study of the $e^+ e^- \rightarrow Z \gamma \gamma \rightarrow q \text{ anti-}q \gamma \gamma$ process at LEP, Phys. Lett. **B505**, 47–58 (2001).
- [72] M. Acciarri et al., Total cross section in $\gamma \gamma$ collisions at LEP, Phys. Lett. **B519**, 33–45 (2001).
- [73] M. Acciarri et al., Measurement of the charm production cross section in $\gamma \gamma$ collisions at LEP, Phys. Lett. **B514**, 19–28 (2001).
- [74] M. Acciarri et al., $K_0(S) K_0(S)$ final state in two-photon collisions and implications for glueballs, Phys. Lett. **B501**, 173–182 (2001).
- [75] M. Acciarri et al., Higgs candidates in $e^+ e^-$ interactions at $\sqrt{s} = 206.6$ - GeV, Phys. Lett. **B495**, 18–25 (2000).
- [76] M. Acciarri et al., Search for excited leptons in $e^+ e^-$ interactions at $\sqrt{s} = 192$ -GeV - 202 -GeV, Phys. Lett. **B502**, 37–50 (2001).
- [77] M. Acciarri et al., Measurements of the cross sections for open charm and beauty production in $\gamma \gamma$ collisions at $\sqrt{s} = 189$ -GeV - 202 -GeV, Phys. Lett. **B503**, 10–20 (2001).
- [78] M. Acciarri et al., Search for neutral Higgs bosons of the minimal supersymmetric standard model in $e^+ e^-$ interactions at $\sqrt{s} = 192$ -GeV to 202 -GeV, Phys. Lett. **B503**, 21–33 (2001).
- [79] M. Acciarri et al., Search for the standard model Higgs boson in $e^+ e^-$ collisions at \sqrt{s} up to 202 -GeV, Phys. Lett. **B508**, 225–236 (2001).
- [80] M. Acciarri et al., Light resonances in $K_0(S) K^+ \pi^-$ and $\eta \pi^+ \pi^-$ final states in $\gamma \gamma$ collisions at LEP, Phys. Lett. **B501**, 1–11 (2001).
- [81] M. Acciarri et al., Search for R-parity violating decays of supersymmetric particles in $e^+ e^-$ collisions at $\sqrt{s} = 189$ -GeV, Eur. Phys. J. **C19**, 397–414 (2001).
- [82] M. Acciarri et al., Study of Z boson pair production in $e^+ e^-$ interactions at $\sqrt{s} = 192$ -GeV to 202 -GeV, Phys. Lett. **B497**, 23–38 (2001).
- [83] M. Acciarri et al., Search for charged Higgs bosons in $e^+ e^-$ collisions at centre-of-mass energies up to 202 -GeV, Phys. Lett. **B496**, 34–42 (2000).

- [84] M. Acciarri et al., Search for a Higgs boson decaying into two photons in $e^+ e^-$ interactions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B489**, 115–124 (2000).
- [85] M. Acciarri et al., Measurement of the $W^+ W^- \gamma$ cross section and direct limits on anomalous quartic gauge boson couplings at LEP, Phys. Lett. **B490**, 187–195 (2000).
- [86] M. Acciarri et al., Search for anomalous couplings in the Higgs sector at LEP, Phys. Lett. **B489**, 102–114 (2000).
- [87] M. Acciarri et al., Measurement of the W pair production cross section and W decay branching fractions in $e^+ e^-$ interactions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B496**, 19–33 (2000).
- [88] M. Acciarri et al., Determination of γ / Z interference in $e^+ e^-$ annihilation at LEP, Phys. Lett. **B489**, 93–101 (2000).
- [89] M. Acciarri et al., Measurement of Bose-Einstein correlations in $e^+ e^- \rightarrow W^+ W^-$ at \sqrt{s} approx. = 189-GeV, Phys. Lett. **B493**, 233–248 (2000).
- [90] M. Acciarri et al., Search for manifestations of new physics in fermion pair production at LEP, Phys. Lett. **B489**, 81–92 (2000).
- [91] M. Acciarri et al., QCD studies in $e^+ e^-$ annihilation from 30-GeV to 189-GeV, Phys. Lett. **B489**, 65–80 (2000).
- [92] M. Acciarri et al., Production of single W bosons at $\sqrt{s} = 189\text{-GeV}$ and measurement of $W W \gamma$ gauge couplings, Phys. Lett. **B487**, 229–240 (2000).
- [93] M. Acciarri et al., Measurements of the b anti- b production cross section and forward-backward asymmetry at centre-of-mass energies above the Z pole at LEP, Phys. Lett. **B485**, 71–84 (2000).
- [94] M. Acciarri et al., Search for anomalous $Z Z \gamma$ and $Z \gamma \gamma$ couplings in the process $e^+ e^- \rightarrow Z \gamma$ at LEP, Phys. Lett. **B489**, 55–64 (2000).
- [95] M. Acciarri et al., Measurement of the lifetime of the tau lepton, Phys. Lett. **B479**, 67–78 (2000).
- [96] M. Acciarri et al., Search for an invisibly decaying Higgs boson in $e^+ e^-$ collisions at $\sqrt{s} = 183\text{-GeV}$ to 189-GeV, Phys. Lett. **B485**, 85–94 (2000).
- [97] M. Acciarri et al., Measurements of cross sections and forward-backward asymmetries at the Z resonance and determination of electroweak parameters, Eur. Phys. J. **C16**, 1–40 (2000).
- [98] M. Acciarri et al., Inclusive Σ^+ and Σ^0 production in hadronic Z decays, Phys. Lett. **B479**, 79–88 (2000).
- [99] M. Acciarri et al., Measurement of the photon structure function at high Q^{*2} at LEP, Phys. Lett. **B483**, 373–386 (2000).
- [100] M. Acciarri et al., Measurement of the running of the fine-structure constant, Phys. Lett. **B476**, 40–48 (2000).
- [101] M. Acciarri et al., Measurement of the $e^+ e^- \rightarrow Z \gamma \gamma$ cross section and determination of quartic gauge boson couplings at LEP, Phys. Lett. **B478**, 39–49 (2000).

- [102] M. Acciarri et al., Search for charginos with a small mass difference with the lightest supersymmetric particle at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B482**, 31–42 (2000).
- [103] M. Acciarri et al., Measurement of hadron and lepton-pair production at $130\text{-GeV} < \sqrt{s} < 189\text{-GeV}$ at LEP, Phys. Lett. **B479**, 101–117 (2000).
- [104] M. Acciarri et al., Hard-photon production and tests of QED at LEP, Phys. Lett. **B475**, 198–205 (2000).
- [105] M. Acciarri et al., Measurement of the probability of gluon splitting into charmed quarks in hadronic Z decays, Phys. Lett. **B476**, 243–255 (2000).
- [106] M. Acciarri et al., Direct observation of longitudinally polarized W^{+-} bosons, Phys. Lett. **B474**, 194–204 (2000).
- [107] M. Acciarri et al., Search for excited leptons at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B473**, 177–185 (2000).
- [108] M. Acciarri et al., Search for neutral Higgs bosons of the minimal supersymmetric standard model in $e^+ e^-$ interactions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B471**, 321–331 (1999).
- [109] M. Acciarri et al., Search for extra dimensions in boson and fermion pair production in $e^+ e^-$ interactions at LEP, Phys. Lett. **B470**, 281–288 (1999).
- [110] M. Acciarri et al., Search for scalar leptons in $e^+ e^-$ collisions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B471**, 280–292 (1999).
- [111] M. Acciarri et al., Search for charginos and neutralinos in $e^+ e^-$ collisions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B472**, 420–433 (2000).
- [112] M. Acciarri et al., Measurement of triple-gauge-boson couplings of the W boson at LEP, Phys. Lett. **B467**, 171–184 (1999).
- [113] M. Acciarri et al., Single and multi-photon events with missing energy in $e^+ e^-$ collisions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B470**, 268–280 (1999).
- [114] M. Acciarri et al., Searches for scalar quarks in $e^+ e^-$ interactions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B471**, 308–320 (1999).
- [115] M. Acciarri et al., Measurement of the spectroscopy of orbitally excited B mesons at LEP, Phys. Lett. **B465**, 323–334 (1999).
- [116] M. Acciarri et al., Search for low scale gravity effects in $e^+ e^-$ collisions at LEP, Phys. Lett. **B464**, 135–144 (1999).
- [117] M. Acciarri et al., Measurement of $R(b)$ and $\text{Br}(b \rightarrow l \nu X)$ at LEP using double-tag methods, Eur. Phys. J. **C13**, 47–61 (2000).
- [118] M. Acciarri et al., Search for charged Higgs bosons in $e^+ e^-$ collisions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B466**, 71–78 (1999).

- [119] M. Acciarri et al., Study of Z boson pair production in $e^+ e^-$ collisions at LEP at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B465**, 363–375 (1999).
- [120] M. Acciarri et al., Measurement of inclusive D^{*+} production in two-photon collisions at LEP, Phys. Lett. **B467**, 137–146 (1999).
- [121] M. Acciarri et al., Search for heavy neutral and charged leptons in $e^+ e^-$ annihilation at $\sqrt{s} = 183\text{-GeV}$ and 189-GeV , Phys. Lett. **B462**, 354–364 (1999).
- [122] M. Acciarri et al., Search for standard model Higgs boson in $e^+ e^-$ interactions at $\sqrt{s} = 189\text{-GeV}$, Phys. Lett. **B461**, 376–386 (1999).
- [123] M. Acciarri et al., Search for heavy isosinglet neutrinos in $e^+ e^-$ annihilation at $130\text{-GeV} < \sqrt{s} < 189\text{-GeV}$, Phys. Lett. **B461**, 397–404 (1999).
- [124] M. Acciarri et al., Formation of the η/c in two-photon collisions at LEP, Phys. Lett. **B461**, 155–166 (1999).
- [125] M. Acciarri et al., Measurement of an elongation of the pion source in Z decays, Phys. Lett. **B458**, 517–528 (1999).
- [126] M. Acciarri et al., Measurement of mass and width of the W boson at LEP, Phys. Lett. **B454**, 386–398 (1999).
- [127] M. Acciarri et al., Search for scalar leptons in $e^+ e^-$ collisions at $\sqrt{s} = 183\text{-GeV}$, Phys. Lett. **B456**, 283–296 (1999).
- [128] M. Acciarri et al., Search for R-parity violating chargino and neutralino decays in $e^+ e^-$ collisions up to $\sqrt{s} = 183\text{-GeV}$, Phys. Lett. **B459**, 354–366 (1999).
- [129] M. Acciarri et al., Study of neutral-current four-fermion and Z Z production in $e^+ e^-$ collisions at $\sqrt{s} = 183\text{-GeV}$, Phys. Lett. **B450**, 281–293 (1999).
- [130] M. Acciarri et al., Measurement of the cross-section for the process $\gamma^* \gamma^* \rightarrow \text{hadrons}$ at LEP, Phys. Lett. **B453**, 333–342 (1999).
- [131] M. Acciarri et al., Heavy quarkonium production in Z decays, Phys. Lett. **B453**, 94–106 (1999).
- [132] M. Acciarri et al., Inclusive charm production in two-photon collisions at LEP, Phys. Lett. **B453**, 83–93 (1999).
- [133] M. Acciarri et al., χ/c_2 formation in two-photon collisions at LEP, Phys. Lett. **B453**, 73–82 (1999).
- [134] M. Acciarri et al., Measurement of the $e^+ e^- \rightarrow Z \rightarrow b \text{ anti-}b$ forward- backward asymmetry and the B_0 anti- B_0 mixing parameter using prompt leptons, Phys. Lett. **B448**, 152–162 (1999).
- [135] M. Acciarri et al., The Q^{*2} evolution of the hadronic photon structure function $F_2(\gamma)$ at LEP, Phys. Lett. **B447**, 147–156 (1999).
- [136] M. Acciarri et al., QCD results from studies of hadronic events produced in $e^+ e^-$ annihilations at $\sqrt{s} = 183\text{-GeV}$, Phys. Lett. **B444**, 569–582 (1998).

- [137] M. Acciarri et al., Search for charged Higgs bosons in $e^+ e^-$ collisions at centre-of-mass energies between 130-GeV and 183-GeV, Phys. Lett. **B446**, 368–377 (1999).
- [138] M. Acciarri et al., Single and multi-photon events with missing energy in $e^+ e^-$ collisions at $\sqrt{s} = 183$ -GeV, Phys. Lett. **B444**, 503–515 (1998).
- [139] M. Acciarri et al., Upper limit on the lifetime difference of short- and long- lived B/s_0 mesons, Phys. Lett. **B438**, 417–429 (1998).
- [140] M. Acciarri et al., Measurement of the effective weak mixing angle by jet- charge asymmetry in hadronic decays of the Z boson, Phys. Lett. **B439**, 225–236 (1998).
- [141] M. Acciarri et al., Searches for scalar top and scalar bottom quarks in $e^+ e^-$ interactions at 161-GeV $\sqrt{s} = 183$ -GeV, Phys. Lett. **B445**, 428–438 (1999).
- [142] M. Acciarri et al., Measurement of W pair cross sections in $e^+ e^-$ interactions at $\sqrt{s} = 183$ -GeV and W decay branching fractions, Phys. Lett. **B436**, 437–452 (1998).
- [143] M. Acciarri et al., Measurement of the Michel parameters and the average tau- neutrino helicity from tau decays at LEP, Phys. Lett. **B438**, 405–416 (1998).
- [144] M. Acciarri et al., Study of anomalous $Z Z \gamma$ and $Z \gamma \gamma$ couplings at LEP, Phys. Lett. **B436**, 187–198 (1998).
- [145] M. Acciarri et al., Measurement of the inclusive charmless semileptonic branching fraction of beauty hadrons and a determination of $|V_{ub}|$ at LEP, Phys. Lett. **B436**, 174–186 (1998).
- [146] M. Acciarri et al., Study of the hadronic photon structure function $F_2(\gamma)$ at LEP, Phys. Lett. **B436**, 403–416 (1998).
- [147] M. Acciarri et al., Production of single W bosons in $e^+ e^-$ interactions at 130- GeV $\sqrt{s} = 183$ -GeV and limits on anomalous $W W \gamma$ couplings, Phys. Lett. **B436**, 417–427 (1998).
- [148] M. Acciarri et al., Test of CP invariance in $Z \rightarrow \mu^+ \mu^- \gamma$ decay, Phys. Lett. **B436**, 428–436 (1998).
- [149] M. Acciarri et al., Measurement of radiative Bhabha and quasi-real Compton scattering, Phys. Lett. **B439**, 183–196 (1998).
- [150] M. Acciarri et al., Search for neutral Higgs bosons of the minimal supersymmetric standard model in $e^+ e^-$ interactions at $\sqrt{s} = 130$ -GeV to 183-GeV, Phys. Lett. **B436**, 389–402 (1998).
- [151] M. Acciarri et al., Photon structure functions and azimuthal correlations of lepton pairs in tagged $\gamma \gamma$ collisions, Phys. Lett. **B438**, 363–378 (1998).
- [152] M. Acciarri et al., Search for the standard model Higgs boson in $e^+ e^-$ interactions at $\sqrt{s} = 183$ -GeV, Phys. Lett. **B431**, 437–450 (1998).
- [153] M. Acciarri et al., Measurement of the anomalous magnetic and electric dipole moments of the tau lepton, Phys. Lett. **B434**, 169–179 (1998).

- [154] M. Acciarri et al., Measurement of the weak dipole moments of the tau lepton, Phys. Lett. **B426**, 207–216 (1998).
- [155] M. Acciarri et al., Angular multiplicity fluctuations in hadronic Z decays and comparison to QCD models and analytical calculations, Phys. Lett. **B428**, 186–196 (1998).
- [156] M. Acciarri et al., Measurement of the B/d0 - anti-B/d0 oscillation frequency, Eur. Phys. J. **C5**, 195–203 (1998).
- [157] M. Acciarri et al., Determination of the number of light neutrino species from single photon production at LEP, Phys. Lett. **B431**, 199–208 (1998).
- [158] M. Acciarri et al., Measurement of tau polarisation at LEP, Phys. Lett. **B429**, 387–398 (1998).
- [159] M. Acciarri et al., Search for new physics phenomena in fermion pair production at LEP, Phys. Lett. **B433**, 163–175 (1998).
- [160] M. Acciarri et al., Local multiplicity fluctuations in hadronic Z decay, Phys. Lett. **B429**, 375–386 (1998).
- [161] M. Acciarri et al., Measurement of the average lifetime of b-hadrons in Z decays, Phys. Lett. **B416**, 220–232 (1998).
- [162] M. Acciarri et al., Search for scalar leptons, charginos and neutralinos in e+ e- collisions at $\sqrt{s} = 161\text{-GeV}$ to 172-GeV , Eur. Phys. J. **C4**, 207–219 (1998).
- [163] M. Acciarri et al., Neutral-current four-fermion production in e+ e- interactions at 130-GeV $\sqrt{s} = 172\text{-GeV}$, Phys. Lett. **B413**, 191–200 (1997).
- [164] M. Acciarri et al., Measurement of $\eta'(958)$ formation in two-photon collisions at LEP1, Phys. Lett. **B418**, 399–410 (1998).
- [165] M. Acciarri et al., Single and multi-photon events with missing energy in e+ e- collisions at $161\text{-GeV} < \sqrt{s} < 172\text{-GeV}$, Phys. Lett. **B415**, 299–310 (1997).
- [166] M. Acciarri et al., Hard-photon production at $\sqrt{s} = 161\text{-GeV}$ and 172-GeV at LEP, Phys. Lett. **B413**, 159–166 (1997).
- [167] M. Acciarri et al., Upsilon production in Z decays, Phys. Lett. **B413**, 167–175 (1997).
- [168] M. Acciarri et al., Search for new physics in energetic single photon production in e+ e- annihilation at the Z resonance, Phys. Lett. **B412**, 201–209 (1997).
- [169] M. Acciarri et al., Search for the standard model Higgs boson in e+ e- interactions at 161-GeV $\sqrt{s} = 172\text{-GeV}$, Phys. Lett. **B411**, 373–386 (1997).
- [170] M. Acciarri et al., Missing mass spectra in hadronic events from e+ e- collisions at $\sqrt{s} = 161\text{-GeV}$ to 172-GeV and limits on invisible Higgs decays, Phys. Lett. **B418**, 389–398 (1998).
- [171] M. Acciarri et al., Measurements of mass, width and gauge couplings of the W boson at LEP, Phys. Lett. **B413**, 176–190 (1997).

- [172] M. Acciarri et al., Search for R-parity breaking sneutrino exchange at LEP, Phys. Lett. **B414**, 373–381 (1997).
- [173] M. Acciarri et al., Measurement of W pair cross sections in e+ e- interactions at $\sqrt{s} = 172$ -GeV and W decay branching fractions, Phys. Lett. **B407**, 419–431 (1997).
- [174] M. Acciarri et al., Resonance formation in the pi+ pi- pi0 final state in two- photon collisions, Phys. Lett. **B413**, 147–158 (1997).
- [175] M. Acciarri et al., Study of hadronic events and measurements of $\alpha(s)$ between 30-GeV and 91-GeV, Phys. Lett. **B411**, 339–353 (1997).
- [176] M. Acciarri et al., Search for heavy neutral and charged leptons in e+ e- annihilation at $\sqrt{s} = 161$ -GeV and $\sqrt{s} = 172$ - GeV, Phys. Lett. **B412**, 189–200 (1997).
- [177] M. Acciarri et al., Cross section of hadron production in gamma gamma collisions at LEP, Phys. Lett. **B408**, 450–464 (1997).
- [178] M. Acciarri et al., K0(S) and Lambda production in quark and gluon jets at LEP, Phys. Lett. **B407**, 389–401 (1997).
- [179] M. Acciarri et al., Measurement of hadron and lepton-pair production at $161\text{-GeV} < \sqrt{s} < 172\text{-GeV}$ at LEP, Phys. Lett. **B407**, 361–376 (1997).
- [180] M. Acciarri et al., Search for anomalous four-jet events in e+ e- annihilation at $\sqrt{s} = 130$ -GeV to 172-GeV, Phys. Lett. **B411**, 330–338 (1997).
- [181] M. Acciarri et al., QCD studies and determination of $\alpha(s)$ in e+ e- collisions at $\sqrt{s} = 161$ -GeV and 172-GeV, Phys. Lett. **B404**, 390–402 (1997).
- [182] M. Acciarri et al., Production of e, mu and tau pairs in untagged two-photon collisions at LEP, Phys. Lett. **B407**, 341–350 (1997).
- [183] M. Acciarri et al., Inclusive J/psi, psi' and chi/c production in hadronic Z decays, Phys. Lett. **B407**, 351–360 (1997).
- [184] M. Acciarri et al., Production of single W bosons at LEP, Phys. Lett. **B403**, 168–176 (1997).
- [185] M. Acciarri et al., Pair-production of W bosons in e+ e- interactions at $\sqrt{s} = 161$ -GeV, Phys. Lett. **B398**, 223–238 (1997).
- [186] M. Acciarri et al., Search for excited leptons in e+ e- annihilation at $\sqrt{s} = 161$ -GeV, Phys. Lett. **B401**, 139–149 (1997).
- [187] M. Acciarri et al., Measurement of $D/s^- \rightarrow \tau^- \text{anti-}\nu/\tau$ and a new limit for $B^- \rightarrow \tau^- \text{anti-}\nu/\tau$, Phys. Lett. **B396**, 327–337 (1997).
- [188] M. Acciarri et al., Study of the weak charged hadronic current in b decays, Phys. Lett. **B393**, 477–486 (1997).
- [189] M. Acciarri et al., Measurement of inclusive omega and eta' production in hadronic Z decays, Phys. Lett. **B393**, 465–476 (1997).

- [190] M. Acciarri et al., Search for exclusive B decays to J and eta or pi0 with the L3 detector, Phys. Lett. **B391**, 481–490 (1997).
- [191] M. Acciarri et al., Search for neutral B meson decays to two charged leptons, Phys. Lett. **B391**, 474–480 (1997).
- [192] M. Acciarri et al., Measurement of the lifetime of the tau lepton, Phys. Lett. **B389**, 187–196 (1996).
- [193] M. Acciarri et al., Search for neutral Higgs boson production through the process $e^+ e^- \rightarrow Z^* H_0$, Phys. Lett. **B385**, 454–470 (1996).
- [194] M. Acciarri et al., Measurement of the B0/d meson oscillation frequency, Phys. Lett. **B383**, 487–498 (1996).
- [195] M. Acciarri et al., Search for new particles in hadronic events with isolated photons, Phys. Lett. **B388**, 409–418 (1996).
- [196] M. Acciarri et al., Measurement of the branching ratios $b \rightarrow e \nu X$, $\mu \nu X$, $\tau \nu X$ and νX , Z. Phys. **C71**, 379–390 (1996).
- [197] M. Acciarri et al., Search for unstable sequential neutral and charged heavy leptons in $e^+ e^-$ annihilation at $\sqrt{s} = 130\text{-GeV}$ and 136-GeV , Phys. Lett. **B377**, 304–312 (1996).
- [198] M. Acciarri et al., Measurement of the Michel parameters and the average tau- neutrino helicity from tau decays in $e^+ e^- \rightarrow \tau^+ \tau^-$, Phys. Lett. **B377**, 313–324 (1996).
- [199] M. Acciarri et al., Observation of multiple hard photon final states at $\sqrt{s} = 130\text{-GeV}$ to 140-GeV at LEP, Phys. Lett. **B384**, 323–332 (1996).
- [200] M. Acciarri et al., Measurement of muon pair production at $50\text{-GeV} < \sqrt{s} \leq 86\text{-GeV}$ at LEP, Phys. Lett. **B374**, 331–340 (1996).
- [201] M. Acciarri et al., Search for supersymmetric particles at $130\text{-GeV} < \sqrt{s} \leq 140\text{-GeV}$ at LEP, Phys. Lett. **B377**, 289–303 (1996).
- [202] M. Acciarri et al., Measurement of eta production in two and three jet events from hadronic Z decays at LEP, Phys. Lett. **B371**, 126–136 (1996).
- [203] M. Acciarri et al., Search for excited leptons in $e^+ e^-$ annihilation at $\sqrt{s} = 130\text{-GeV} - 140\text{-GeV}$, Phys. Lett. **B370**, 211–221 (1996).
- [204] M. Acciarri et al., Measurement of hadron and lepton pair production at $130\text{-GeV} < \sqrt{s} < 140\text{-GeV}$ at LEP, Phys. Lett. **B370**, 195–210 (1996).
- [205] M. Acciarri et al., Study of the structure of hadronic events and determination of α_s at $\sqrt{s} = 130\text{-GeV}$ and 136-GeV , Phys. Lett. **B371**, 137–148 (1996).
- [206] M. Acciarri et al., Search for the decays $B(d)0 \rightarrow \gamma \gamma$ and $B(s)0 \rightarrow \gamma \gamma$, Phys. Lett. **B363**, 137–144 (1995).

- [207] M. Acciarri et al., Study of the $K(s)0 K(s)0$ final state in two photon collisions, Phys. Lett. **B363**, 118–126 (1995).
- [208] M. Acciarri et al., Search for neutral charmless B decays at LEP, Phys. Lett. **B363**, 127–136 (1995).
- [209] M. Acciarri et al., Evidence for gluon interference in hadronic Z decays, Phys. Lett. **B353**, 145–154 (1995).
- [210] M. Acciarri et al., One prong tau decays with neutral kaons, Phys. Lett. **B352**, 487–497 (1995).
- [211] M. Acciarri et al., Tests of QED at LEP energies using $e^+ e^- \rightarrow \gamma \gamma$ (γ) and $e^+ e^- \rightarrow \text{lepton}^+ \text{lepton}^- \gamma \gamma$, Phys. Lett. **B353**, 136–144 (1995).
- [212] M. Acciarri et al., Search for neutralinos in Z decays, Phys. Lett. **B350**, 109–119 (1995).
- [213] M. Acciarri et al., Measurement of the weak charged current structure in semileptonic b hadron decays at the Z peak, Phys. Lett. **B351**, 375–386 (1995).
- [214] M. Acciarri et al., Measurement of energetic single photon production at LEP, Phys. Lett. **B346**, 190–198 (1995).
- [215] M. Acciarri et al., Measurement of exclusive branching fractions of hadronic one space prong tau decays, Phys. Lett. **B345**, 93–102 (1995).
- [216] M. Acciarri et al., Search for anomalous $Z \rightarrow \gamma \gamma \gamma$ events at LEP, Phys. Lett. **B345**, 609–616 (1995).
- [217] M. Acciarri et al., Energy and particle flow in three jet and radiative two jet events from hadronic Z decays, Phys. Lett. **B345**, 74–84 (1995).
- [218] M. Acciarri et al., A Measurement of tau polarization at LEP, Phys. Lett. **B341**, 245–256 (1994).
- [219] M. Acciarri et al., B^* production in Z decays at LEP, Phys. Lett. **B345**, 589–597 (1995).
- [220] M. Acciarri et al., Measurement of the $B^0 - \text{anti-}B^0$ mixing parameter and the $Z \rightarrow b \text{ anti-}b$ forward - backward asymmetry, Phys. Lett. **B335**, 542–553 (1994).
- [221] M. Acciarri et al., Measurement of the inclusive $B \rightarrow \text{tau-neutrino } X$ branching ratio, Phys. Lett. **B332**, 201–208 (1994).
- [222] M. Acciarri et al., Measurement of cross-sections and leptonic forward - backward asymmetries at the z pole and determination of electroweak parameters, Z. Phys. **C62**, 551–576 (1994).
- [223] M. Acciarri et al., Measurement of inclusive production of neutral hadrons from Z decays, Phys. Lett. **B328**, 223–233 (1994).
- [224] A. Adam et al., A Study of four fermion processes at LEP, Phys. Lett. **B321**, 283–294 (1994).
- [225] O. Adriani et al., Measurement of $\eta(c)$ production in untagged two photon collisions at LEP, Phys. Lett. **B318**, 575–582 (1993).
- [226] O. Adriani et al., An S matrix analysis of the Z resonance, Phys. Lett. **B315**, 494–502 (1993).

- [227] O. Adriani et al., Search for lepton flavor violation in Z decays, Phys. Lett. **B316**, 427–434 (1993).
- [228] O. Adriani et al., Measurement of the average lifetime of b hadrons, Phys. Lett. **B317**, 474–484 (1993).
- [229] O. Adriani et al., Chi(c) production in hadronic Z decays, Phys. Lett. **B317**, 467–473 (1993).
- [230] O. Adriani et al., Inclusive search for the charmless radiative decay of the b quark ($b \rightarrow s \gamma$), Phys. Lett. **B317**, 637–646 (1993).
- [231] O. Adriani et al., Measurement of $\Gamma(b \text{ anti-}b) / \Gamma(b \text{ had})$ from hadronic decays of the Z, Phys. Lett. **B307**, 237–246 (1993).
- [232] O. Adriani et al., Determination of the effective electroweak mixing angle from Z decay, Phys. Lett. **B309**, 451–462 (1993).
- [233] O. Adriani et al., Search for narrow vector resonances in the Z mass range, Phys. Lett. **B313**, 326–332 (1993).
- [234] O. Adriani et al., Search for a Z-prime at the Z resonance, Phys. Lett. **B306**, 187–196 (1993).
- [235] O. Adriani et al., Results from the L3 experiment at LEP, Phys. Rept. **236**, 1–146 (1993).
- [236] O. Adriani et al., A Search for the neutral Higgs boson at LEP, Phys. Lett. **B303**, 391–399 (1993).
- [237] O. Adriani et al., Determination of quark electroweak couplings from direct photon production in hadronic Z decays, Phys. Lett. **B301**, 136–144 (1993).
- [238] O. Adriani et al., Search for anomalous production of single photon events in $e^+ e^-$ annihilations at the Z resonance, Phys. Lett. **B297**, 469–476 (1992).
- [239] O. Adriani et al., Search for isosinglet neutral heavy leptons in Z^0 decays, Phys. Lett. **B295**, 371–382 (1992).
- [240] O. Adriani et al., High mass photon pairs in $\text{lepton}^+ \text{lepton}^- \gamma \gamma$ events at LEP, Phys. Lett. **B295**, 337–346 (1992).
- [241] O. Adriani et al., Search for nonminimal Higgs bosons in Z^0 decays, Z. Phys. **C57**, 355–382 (1993).
- [242] O. Adriani et al., Determination of the number of light neutrino species, Phys. Lett. **B292**, 463–471 (1992).
- [243] O. Adriani et al., A Measurement of tau polarization in Z^0 decays, Phys. Lett. **B294**, 466–478 (1992).
- [244] O. Adriani et al., Searches for nonminimal Higgs bosons in Z^0 decays, Phys. Lett. **B294**, 457–465 (1992).
- [245] O. Adriani et al., Measurement of the $e^+ e^- \rightarrow b \text{ anti-}b$ and $e^+ e^- \rightarrow c \text{ anti-}c$ forward backward asymmetries at the Z^0 resonance, Phys. Lett. **B292**, 454–462 (1992).

- [246] O. Adriani et al., Isolated hard photon emission in hadronic Z0 decays, Phys. Lett. **B292**, 472–484 (1992).
- [247] O. Adriani et al., Inclusive J production in Z0 decays, Phys. Lett. **B288**, 412–420 (1992).
- [248] B. Adeva et al., An Improved measurement of B0 - anti-B0 mixing in Z0 decays, Phys. Lett. **B288**, 395–403 (1992).
- [249] O. Adriani et al., Measurement of inclusive eta production in hadronic decays of the Z0, Phys. Lett. **B286**, 403–412 (1992).
- [250] O. Adriani et al., A Test of quantum electrodynamics in the reaction $e^+ e^- \rightarrow \gamma \gamma$ ($\gamma \gamma$), Phys. Lett. **B288**, 404–411 (1992).
- [251] O. Adriani et al., Determination of α_s from hadronic event shapes measured on the Z0 resonance, Phys. Lett. **B284**, 471–481 (1992).
- [252] B. Adeva et al., Search for the neutral Higgs boson at LEP, Phys. Lett. **B283**, 454–464 (1992).
- [253] B. Adeva et al., Studies of hadronic event structure and comparisons with QCD models at the Z0 resonance, Z. Phys. **C55**, 39–62 (1992).
- [254] B. Adeva et al., A Direct determination of the number of light neutrino families from $e^+ e^- \rightarrow \nu \bar{\nu} \gamma$ at LEP, Phys. Lett. **B275**, 209–221 (1992).
- [255] B. Adeva et al., Measurement of the strong coupling constant α_s for bottom quarks at the Z0 resonance, Phys. Lett. **B271**, 461–467 (1991).
- [256] B. Adeva et al., Search for lepton flavor violation in Z0 decays, Phys. Lett. **B271**, 453–460 (1991).
- [257] B. Adeva et al., Measurement of the lifetime of B hadrons and a determination of $|V_{cb}|$, Phys. Lett. **B270**, 111–122 (1991).
- [258] B. Adeva et al., Decay properties of tau leptons measured at the Z0 resonance, Phys. Lett. **B265**, 451–461 (1991).
- [259] B. Adeva et al., A Test of QCD based on three jet events from Z0 decays, Phys. Lett. **B263**, 551–562 (1991).
- [260] B. Adeva et al., Search for leptoquarks in Z0 decays, Phys. Lett. **B261**, 169–176 (1991).
- [261] B. Adeva et al., Measurement of the inclusive production of neutral pions and charged particles on the Z0 resonance, Phys. Lett. **B259**, 199–208 (1991).
- [262] B. Adeva et al., Measurements of $Z0 \rightarrow b \bar{b}$ decays and the semileptonic branching ratio $Br(b \rightarrow \text{lepton } X)$, Phys. Lett. **B261**, 177–187 (1991).
- [263] B. Adeva et al., Measurement of electroweak parameters from hadronic and leptonic decays of the Z0, Z. Phys. **C51**, 179–204 (1991).
- [264] B. Adeva et al., Search for narrow high mass resonances in radiative decays of the Z0, Phys. Lett. **B262**, 155–162 (1991).

- [265] B. Adeva et al., Determination of α_s from energy-energy correlations measured on the Z^0 resonance, Phys. Lett. **B257**, 469–478 (1991).
- [266] B. Adeva et al., Search for the neutral Higgs boson, Phys. Lett. **B257**, 450–458 (1991).
- [267] B. Adeva et al., A Measurement of the $Z^0 \rightarrow b$ anti- b forward - backward asymmetry, Phys. Lett. **B252**, 713–721 (1990).
- [268] B. Adeva et al., A Measurement of B^0 - anti- B^0 mixing in Z^0 decays, Phys. Lett. **B252**, 703–712 (1990).
- [269] B. Adeva et al., Search for excited neutrinos from Z^0 decays, Phys. Lett. **B252**, 525–532 (1990).
- [270] B. Adeva et al., Search for the charged Higgs boson in Z^0 decay, Phys. Lett. **B252**, 511–517 (1990).
- [271] B. Adeva et al., Test of QED in $e^+ e^- \rightarrow \gamma \gamma$ at LEP, Phys. Lett. **B250**, 199–204 (1990).
- [272] B. Adeva et al., Search for excited taus from Z^0 decay, Phys. Lett. **B250**, 205–211 (1990).
- [273] B. Adeva et al., Search for the neutral Higgs bosons of the minimal supersymmetric standard model from Z^0 decays, Phys. Lett. **B251**, 311–320 (1990).
- [274] B. Adeva et al., A Search for heavy charged and neutral leptons from Z^0 decays, Phys. Lett. **B251**, 321–330 (1990).
- [275] B. Adeva et al., A Determination of electroweak parameters from Z^0 decays into charged leptons, Phys. Lett. **B250**, 183–192 (1990).